

## NCTN Leukemia Trials Portfolio (Open as of 2/15/2025)

Each far right box includes the NCTN protocol number with a hyperlink to the associated ClinicalTrials.gov webpage. Click on it to view the protocol title and study information.



## MyeloMATCH and its sub-trials:



## NCTN Leukemia Trials (Open as of 2/15/2025)

<b>Protocol Number</b>	Phase	Protocol Title
		A Phase III Trial to Evaluate the Efficacy of the Addition of Inotuzumab Ozogamicin (a Conjugated Anti-CD22
		Monoclonal Antibody) to Frontline Therapy in Young Adults (Ages 18-39 Years) with Newly Diagnosed
A041501	Ш	Precursor B-Cell ALL
		A Phase II Study of Inotuzumab Ozogamicin Followed by Blinatumomab for Ph-Negative, CD22-Positive B-
		Lineage Acute Lymphoblastic Leukemia in Newly Diagnosed Older Adults or Adults with Relapsed or Refractory
A041703	II	Disease
		A Randomized Phase II Study Comparing Inotuzumab Plus Chemotherapy Versus Standard Chemotherapy in
A042001	II	Older Adults with Philadelphia-Chromosome-Negative B-Cell Acute Lymphoblastic Leukemia
		A Phase III Randomized Trial of Steroids + Tyrosine Kinase Inhibitor Induction with Chemotherapy or
EA9181	Ш	Blinatumomab for Newly Diagnosed BCR-ABL-Positive Acute Lymphoblastic Leukemia in Adults
		A Phase II Study of Daratumumab-Hyaluronidase for Chemotherapy-Relapsed/Refractory Minimal Residual
EA9213	II	Disease (MRD) in T Cell Acute Lymphoblastic Leukemia (T-ALL)
		MYELOMATCH, Master Screening and Reassessment Protocol (MSRP) for Tier Advancement in the NCI
MYELOMATCH	Other	MyeloMATCH Clinical Trials
		A Measurable Residual Disease (MRD) Focused, Phase II Study of Venetoclax Plus Chemotherapy for Newly
		Diagnosed Younger Patients with Intermediate Risk Acute Myeloid Leukemia: A Tier 1 MYELOMATCH Clinical
MM1YA-CTG01	II	Trial
		A Randomized Phase II Study of Venetoclax and HMA-Based Therapies for the Treatment of Older and Unfit
MM1OA-EA02	п	Adults with Newly Diagnosed FLT3-Mutated Acute Myeloid Leukemia (AML): A MyeloMATCH Treatment Trial
		,
		A Randomized Phase II Study Comparing Cytarabine + Daunorubicin (7 + 3) vs (Daunorubicin and Cytarabine)
		Liposome, Cytarabine + Daunorubicin + Venetoclax, Azacitidine + Venetoclax, and (Daunorubicin and
		Cytarabine) Liposome + Venetoclax in Patients Aged 59 or Younger Who are Considered High-Risk (Adverse)
MM1YA-S01	II	Acute Myeloid Leukemia As Determined by MYELOMATCH; A MYELOMATCH Clinical Trial
		A Phase I/II Study of AKR1C3-Activated Prodrug OBI-3424 (OBI-3424) In Patients with Relapsed/Refractory T-
S1905	II	Cell Acute Lymphoblastic Leukemia (T-ALL)/T-Cell Lymphoblastic Lymphoma (T-LBL)
		Randomized, Phase III Study of Early Intervention with Venetoclax and Obinutuzumab Versus Delayed Therapy
		with Venetoclax and Obinutuzumab in Newly Diagnosed Asymptomatic High-Risk Patients with Chronic
S1925	Ш	Lymphocytic Leukemia/Small Lymphocytic Lymphoma (CLL/SLL): EVOLVE CLL/SLL Study