



NCTN Leukemia Trials Portfolio (Open as of 3/15/2025)

Each far right box includes the NCTN protocol number with a hyperlink to the associated ClinicalTrials.gov webpage. Click on it to view the protocol title and study information.



MyeloMATCH and its sub-trials:



NCTN Leukemia Trials (Open as of 3/15/2025)

Protocol Number	Phase	Protocol Title
A041501	III	A Phase III Trial to Evaluate the Efficacy of the Addition of Inotuzumab Ozogamicin (a Conjugated Anti-CD22 Monoclonal Antibody) to Frontline Therapy in Young Adults (Ages 18-39 Years) with Newly Diagnosed Precursor B-Cell ALL
A041703	II	A Phase II Study of Inotuzumab Ozogamicin Followed by Blinatumomab for Ph-Negative, CD22-Positive B-Lineage Acute Lymphoblastic Leukemia in Newly Diagnosed Older Adults or Adults with Relapsed or Refractory Disease
A042001	II	A Randomized Phase II Study Comparing Inotuzumab Plus Chemotherapy Versus Standard Chemotherapy in Older Adults with Philadelphia-Chromosome-Negative B-Cell Acute Lymphoblastic Leukemia
EA9181	III	A Phase III Randomized Trial of Steroids + Tyrosine Kinase Inhibitor Induction with Chemotherapy or Blinatumomab for Newly Diagnosed BCR-ABL-Positive Acute Lymphoblastic Leukemia in Adults
EA9213	II	A Phase II Study of Daratumumab-Hyaluronidase for Chemotherapy-Relapsed/Refractory Minimal Residual Disease (MRD) in T Cell Acute Lymphoblastic Leukemia (T-ALL)
MYELOMATCH	Other	MYELOMATCH, Master Screening and Reassessment Protocol (MSRP) for Tier Advancement in the NCI MyeloMATCH Clinical Trials
MM1YA-CTG01	II	A Measurable Residual Disease (MRD) Focused, Phase II Study of Venetoclax Plus Chemotherapy for Newly Diagnosed Younger Patients with Intermediate Risk Acute Myeloid Leukemia: A Tier 1 MYELOMATCH Clinical Trial
MM10A-EA02	II	A Randomized Phase II Study of Venetoclax and HMA-Based Therapies for the Treatment of Older and Unfit Adults with Newly Diagnosed FLT3-Mutated Acute Myeloid Leukemia (AML): A MyeloMATCH Treatment Trial
MM1YA-S01	II	A Randomized Phase II Study Comparing Cytarabine + Daunorubicin (7 + 3) vs (Daunorubicin and Cytarabine) Liposome, Cytarabine + Daunorubicin + Venetoclax, Azacitidine + Venetoclax, and (Daunorubicin and Cytarabine) Liposome + Venetoclax in Patients Aged 59 or Younger Who are Considered High-Risk (Adverse) Acute Myeloid Leukemia As Determined by MYELOMATCH; A MYELOMATCH Clinical Trial
S1905	II	A Phase I/II Study of AKR1C3-Activated Prodrug OBI-3424 (OBI-3424) In Patients with Relapsed/Refractory T-Cell Acute Lymphoblastic Leukemia (T-ALL)/T-Cell Lymphoblastic Lymphoma (T-LBL)
S1925	III	Randomized, Phase III Study of Early Intervention with Venetoclax and Obinutuzumab Versus Delayed Therapy with Venetoclax and Obinutuzumab in Newly Diagnosed Asymptomatic High-Risk Patients with Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma (CLL/SLL): EVOLVE CLL/SLL Study